

(tangent (hyper-)plane)

1. Given $f(x, y) = 3y^2 - 2x^2 + x$ and a point $T = [2; -1; ?]$.
 - a) Compute P.D. of the function in a point $[2; -1]$.
 - b) Find an equation of the tangent plane (τ) to the graph of the function at the point T .
2. Find an equation of the plane (τ) tangent to the graph of $f(x, y) = x \sin(x + y)$ at a point $T = [-1; 1; ?]$. Find also an equation of a line (ν) normal to the graph of f at point T .
3. a) Find an equation of the plane tangent to the graph of $f(x, y) = \ln(x + y)$ at a point $[1; 0; ?]$.
b) Use the result to approximate the functional value $f(A_1)$ in a point $A_1 = [1.1; 0.1]$.
4. Given $f(x, y) = 2x^2 - y^2$ and a plane $\sigma : 8x - 6y - z + 12 = 0$.
 - a) Find a plane (τ) tangent to the graph of f and parallel to the plane σ .
 - b) Find a line (ν) normal to the graph of f and normal to the plane σ .
5. Find an equation of the hyper-plane (τ) tangent to the graph of $f(x, y, z) = \ln(x^2 - y + 3z)$ at a point $T = [2; 1; 1; ?]$.
6. Given $f(x, y, z) = \ln(z + \sqrt{9 - x^2 - y^2})$,
 - a) Find Domain of definition of f and sketch it (at least in 2 cuts).
 - b) Find an equation of the hyper-plane (τ) tangent to the graph of f at a point $T = [0; 0; 1; ?]$.

Gradient and directional derivative

7. Given $f(x, y) = \sqrt{1 - x^2} - \sqrt{1 - y^2}$,
 - a) find Domain of definition of f and sketch it.
 - b) Where is the function f differentiable? (Find the domain of differentiability.)
 - c) Compute gradient of the function in a point $A = [1/2; 0]$
8. Given $f(x, y) = \frac{\sqrt{y-x^2}}{1-x^2}$ and a point $A = [0; 1]$,
 - a) find the Domain of definition of f and sketch it.
 - b) Where is the function f differentiable? (Find the domain of differentiability.)
 - c) Determine the direction in which the graph of the function is increasing the most at point A .
9. Given $f(x, y, z) = \sin xz + x + y - \frac{z}{y}$ and a point $A = [2; 1; 0]$,
Determine the direction of maximal decrease of the function f at the point A .
10. Given $f(x, y) = x^2 + 2xy - 3y^2$ and a point $A = [1; 1]$,
 - a) compute the (directional) derivative of f at point A in direction given by vector $\vec{s} = (3; 4)$.
 - b) Describe the behavior of the function in this direction.
 - c) Compute the derivative of f at point A in the direction given by the vector $\vec{t} = \frac{1}{\sqrt{2}}(1; 1)$.
What can you say about the function in this direction at the point A ?
11. Given $f(x, y) = \cos xy + e^{xy}$ and a point $A = [1; 0]$,
 - a) determine the direction \vec{s} of maximal increase of the function f at a point A .
 - b) Compute the (directional) derivative of f at point A in the direction given by a vector \vec{s} .
 - c) Compute the derivative of f at point A in the direction given by a vector $\vec{t} = (1; 2)$. What can you say about the function in this direction?
12. Given $f(x, y) = \sqrt{9 - x - y^2}$ and a point $A = [1; -2]$,
 - a) compute gradient of the function at point A .
 - b) Find the direction vector \vec{u} in which the function doesn't change its value.